

## Atypical lipid keratopathy

Kammoun Sonda; Rekik Mona\*; Gargouri Salma; Trigui Amira

Department of ophthalmology, Habib Bourguiba Hospital, Sfax, Tunisia.

**Received Date** : July 28, 2023  
**Accepted Date** : September 11, 2023  
**Published Date** : September 18, 2023  
**Archived** : [www.jcmimagescasereports.org](http://www.jcmimagescasereports.org)  
**Copyright** : © Mona Rekik 2023

**\*Corresponding Author:** Mona Rekik, Department of ophthalmology, Habib Bourguiba Hospital, Sfax, Tunisia.  
Tel: +00-216-24300099;  
Email: rekikmona@yahoo.fr

### Clinical Image


A 26-year-old patient, presented with a bilateral corneal clouding that had been slowly progressive over the last 2 years. He was a sports trainer having consumed protein for bodybuilding. Visual acuity was 6/10 in the Right Eye (RE) and 1/20 in the Left Eye (LE).

Slit lamp examination showed deep yellow corneal deposits with stromal neovascularization (Figure 1). No other ocular abnormalities were found. The anterior segment optical coherence tomography revealed a hyper-reflective area in the deep stroma (Figure 2). Confocal microscopy showed hyper-reflective structures in deep stroma (Figure 3). Evidence of hyperlipidemia was not apparent. No evidence was found for

a systemic or an infectious disease. The patient was diagnosed with lipid keratopathy secondary to anabolics. This case describes a very rare condition and represents the first case of lipid keratopathy secondary to anabolics in a sports trainer. To the best of our knowledge, this is the first report of corneal neovascularization in a patient using anabolics reported to date.


Lipid keratopathy is a rare disease. It has a primary and a secondary form, the secondary being the more common entity because of leakage from lipids out of newly formed corneal vessels after inflammation. Oxidative stress and physical exercise increase surface inflammation and limbitis and promote corneal neovascularization which leads lipid keratopathy.

**Declarations of interest:** None.




**Figure 1:** Corneal deposits in the right and the left eye with stromal neovascularization in the left eye.

**Citation:** Mona Rekik. Atypical lipid keratopathy. J Clin Med Img Case Rep. 2023; 3(5): 1550.



**Figure 2:** Anterior segment optical coherence tomography revealing a hyper-reflective area in the deep stroma.



**Figure 3:** Confocal microscopy showing increased number of Keratocytes with pathological activity proving inflammatory state.